

ON THE VIRIAL THEOREM FOR THE RELATIVISTIC OPERATOR OF BROWN AND RAVENHALL, AND THE ABSENCE OF EMBEDDED EIGENVALUES

A. A. BALINSKY AND W. D. EVANS

ABSTRACT. A virial theorem is established for the operator proposed by Brown and Ravenhall as a model for relativistic one-electron atoms. As a consequence, it is proved that the operator has no eigenvalues greater than $\max(mc^2, 2\alpha Z - \frac{1}{2})$, where α is the fine structure constant, for all values of the nuclear charge Z below the critical value Z_c : in particular there are no eigenvalues embedded in the essential spectrum when $Z \leq 3/4\alpha$. Implications for the operators in the partial wave decomposition are also described.

1. INTRODUCTION

The formal operator proposed by Brown and Ravenhall in [1] to include relativistic effects in the description of an electron in the field of its nucleus is of the form

$$\mathbf{B} := \Lambda_+ \left(D_0 - \frac{e^2 Z}{|\cdot|} \right) \Lambda_+. \quad (1.1)$$

In (1.1), the notation is as follows (see [3]):

- D_0 is the free Dirac operator

$$D_0 = c\boldsymbol{\alpha} \cdot \frac{\hbar}{i} \nabla + mc^2\beta \equiv \sum_{j=1}^3 c \frac{\hbar}{i} \alpha_j \frac{\partial}{\partial x_j} + mc^2\beta,$$

where $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$ and β are the Dirac matrices given by

$$\alpha_j = \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1_2 & 0_2 \\ 0_2 & -1_2 \end{pmatrix}$$

with $\mathbf{0}_2, \mathbf{1}_2$ the zero and unit 2×2 matrices respectively and σ_j the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- Λ_+ denotes the projection of $L^2(\mathbb{R}^3) \otimes \mathbb{C}^4$ onto the positive spectral subspace of D_0 , that is $\chi_{(0,\infty)}(D_0)$, where $\chi_{(0,\infty)}$ is the characteristic function of $(0, \infty)$. If we set

$$\widehat{f}(\mathbf{p}) \equiv \mathcal{F}(f)(\mathbf{p}) = \left(\frac{1}{2\pi\hbar} \right)^{3/2} \int_{\mathbb{R}^3} e^{-i \mathbf{x} \cdot \mathbf{p}/\hbar} f(\mathbf{x}) \, d\mathbf{x}$$

Date: February 1, 2008.

1991 Mathematics Subject Classification. Primary: 47G, 81Q10; Secondary: 47N50.

Key words and phrases. Brown-Ravenhall operator, virial theorem, essential spectra, eigenvalues.

for the Fourier transform of f , then it follows that

$$(\Lambda_+ f)^\wedge(\mathbf{p}) = \Lambda_+(\mathbf{p}) \widehat{f}(\mathbf{p}),$$

where

$$\Lambda_+(\mathbf{p}) = \frac{1}{2} + \frac{c\boldsymbol{\alpha} \cdot \mathbf{p} + mc^2\beta}{2\mathbf{e}(p)}, \quad \mathbf{e}(p) = \sqrt{c^2p^2 + m^2c^4} \quad (1.2)$$

with $p = |\mathbf{p}|$.

- $2\pi\hbar$ is Planck's constant, c the velocity of light, m the electron mass, $-e$ the electron charge, and Z the nuclear charge.

The underlying Hilbert space in which $\mathbf{B}$ acts is

$$\mathcal{H} = \Lambda_+(\mathbb{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^4). \quad (1.3)$$

The Fourier transform of any spinor ψ in the positive spectral subspace of D_0 can be written

$$\widehat{\psi}(\mathbf{p}) = \frac{1}{\mathbf{n}(p)} \begin{pmatrix} [\mathbf{e}(p) + \mathbf{e}(0)] u(\mathbf{p}) \\ c(\mathbf{p} \cdot \boldsymbol{\sigma}) u(\mathbf{p}) \end{pmatrix}, \quad (1.4)$$

where $u \in \mathbb{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^2$, a Pauli spinor, and $\mathbf{n}(p) = [2\mathbf{e}(p)(\mathbf{e}(p) + \mathbf{e}(0))]^{1/2}$. Conversely any Dirac spinor of the form (1.4) is in the image of $\mathcal{H}$ under the Fourier transform. We then have formally that, if v is the Pauli spinor related by (1.4) to a Dirac spinor φ ,

$$(\varphi, \mathbf{B}\psi) = \boldsymbol{\beta}[v, u] := \mathbf{e}[v, u] - \gamma \mathbf{k}[v, u], \quad (1.5)$$

where $(\cdot, \cdot)$ is the inner-product on $(\mathbb{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^4)$ (being linear in the second argument),

$$\mathbf{e}[v, u] = \int_{\mathbb{R}^3} \mathbf{e}(p) v(\mathbf{p})^* u(\mathbf{p}) d\mathbf{p}, \quad (1.6)$$

$$\mathbf{k}[v, u] = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} v(\mathbf{p}')^* \mathbf{K}(\mathbf{p}', \mathbf{p}) u(\mathbf{p}) d\mathbf{p}' d\mathbf{p}, \quad (1.7)$$

and

$$\gamma = \frac{\alpha c Z}{2\pi^2}. \quad (1.8)$$

In (1.6)-(1.8), $*$ denotes the Hermitian conjugate, $\alpha = e^2/(\hbar c)$ is Sommerfeld's fine structure constant and the kernel $\mathbf{K}$ in (1.7) is the 2×2 matrix-valued function

$$\mathbf{K}(\mathbf{p}', \mathbf{p}) = \frac{[\mathbf{e}(p') + \mathbf{e}(0)][\mathbf{e}(p) + \mathbf{e}(0)]\mathbf{1}_2 + c^2(\mathbf{p}' \cdot \boldsymbol{\sigma})(\mathbf{p} \cdot \boldsymbol{\sigma})}{\mathbf{n}(p') |\mathbf{p} - \mathbf{p}'|^2 \mathbf{n}(p)}. \quad (1.9)$$

The form $\mathbf{e}[\cdot]$ with domain $\mathbb{L}^2(\mathbb{R}^3; \sqrt{1+p^2} d\mathbf{p}) \otimes \mathbb{C}^2$ is closed and non-negative in $\mathbb{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^2$. In [2] and [7] it was proved that for $u \in \mathbb{L}^2(\mathbb{R}^3; \sqrt{1+p^2} d\mathbf{p}) \otimes \mathbb{C}^2$ and $Z \leq Z_c = 2/[(\frac{\pi}{2} + \frac{2}{\pi})\alpha]$,

$$\gamma \mathbf{k}[u, u] \leq \frac{Z}{Z_c} \mathbf{e}[u, u]. \quad (1.10)$$

It had earlier been established in [3] that $\boldsymbol{\beta}$ is bounded below if and only if $Z \leq Z_c$, confirming a prediction of Hardenkopf and Sucher [4] based on numerical considerations.

The strict positivity of β is proved in both [2] [7], a positive lower bound being exhibited in [7] even for $Z = Z_c$. If $Z < Z_c$, it follows from (1.5) and (1.10) that β is a closed positive form with domain $\mathcal{D}(\beta) = L^2(\mathbb{R}^3; \sqrt{1+p^2} d\mathbf{p}) \otimes \mathbb{C}^2$. There is therefore defined a positive self-adjoint operator $\mathbf{b}$ in $L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ satisfying

$$(v, \mathbf{b}u) = \beta[v, u], \quad u \in \mathcal{D}(\mathbf{b}), \quad v \in \mathcal{D}(\beta), \quad (1.11)$$

where the domain $\mathcal{D}(\mathbf{b})$ of $\mathbf{b}$ is dense in its “form domain” $\mathcal{D}(\beta)$. We shall follow [3] and refer to this operator $\mathbf{b}$ as the Brown-Ravenhall operator, it being associated to the original operator $\mathbf{B}$ via (1.5).

If $Z = Z_c$, the operator $\mathbf{b}$ is defined as the Friedrichs operator associated with the closure of the form β restricted to rapidly decreasing Pauli spinors. The description of the operator $\mathbf{b}$ in this case is studied in [8]. We assume in this paper that $Z < Z_c$, so that $\mathbf{b}$ is defined in (1.11). In this case, it follows from [3, Theorem 2] that $\sigma_{\text{ess}}(\mathbf{b}) = [mc^2, \infty)$ and $\sigma_{\text{sc}}(\mathbf{b}) = \emptyset$, where σ_{ess} and σ_{sc} are respectively the essential and singular continuous spectra. We shall establish a virial theorem for $\mathbf{b}$ which will imply that for $Z \leq Z'_c = \frac{3}{4\alpha}$ there are no eigenvalues embedded in $[mc^2, \infty)$, and hence the spectrum in $[mc^2, \infty)$ is absolutely continuous.

The number Z'_c has a spectral significance in [8] concerning the self-adjointness of operators $b_{l,s}$ in the partial wave decomposition of $\mathbf{B}$ and $\mathbf{b}$: in $b_{l,s}$, l denotes the angular momentum channel and s the spin, and in [8] the operators are given the domain $L^2(0, \infty; [\sqrt{1+p^2}] dp)$. Tix proves that for $(l, s) \neq (0, 1/2)$ or $(1, -1/2)$, the operators $b_{l,s}$ are all self-adjoint for $Z < Z_c$, but $b_{0,1/2}$ and $b_{1,-1/2}$ are self-adjoint if $Z < Z'_c$, essentially self-adjoint when $Z = Z'_c$ and are symmetric with a one-parameter family of self-adjoint extensions when $Z'_c < Z \leq Z_c$. We also prove that for all $(l, s) \neq (0, 1/2)$ or $(1, -1/2)$, the $b_{l,s}$ have no eigenvalues in $[mc^2, \infty)$ for $Z < Z_c$, but only for $Z \leq Z'_c$ in the case $(l, s) = (1, -1/2)$. The operator $b_{0,1/2}$ (its Friedrichs extension when $Z \geq Z'_c$) has no eigenvalues in $[mc^2, \infty)$ for the whole range $Z < Z_c$.

2. THE VIRIAL THEOREM

We begin by proving an abstract virial theorem.

Lemma 2.1. *Let $U(a)$, $a \in \mathbb{R}_+$, be a one parameter family of unitary operators on a Hilbert space $\mathcal{H}$ which converges strongly to the identity as $a \rightarrow 1$. Let T be a self-adjoint operator in $\mathcal{H}$ and $T_a = f(a) U(a) T U(a)^{-1}$, where $f(1) = 1$ and $f'(1)$ exists. If $\varphi \in \mathcal{D}(T) \cap \mathcal{D}(T_a)$ is an eigenvector of T corresponding to an eigenvalue λ then*

$$\lim_{a \rightarrow 1} \left(\varphi_a, \left[\frac{T_a - T}{a - 1} \right] \varphi \right) = \lambda f'(1) \|\varphi\|^2, \quad (2.1)$$

where $\varphi_a = U(a)\varphi$.

Proof. From $T\varphi = \lambda\varphi$ we have $T_a\varphi_a = \lambda f(a)\varphi_a$. Hence

$$(\varphi_a, T\varphi) = \lambda(\varphi_a, \varphi)$$

and

$$(T_a \varphi_a, \varphi) = \lambda f(a)(\varphi_a, \varphi).$$

Consequently

$$\left(\varphi_a, \left[\frac{T_a - T}{a - 1} \right] \varphi \right) = \lambda \left[\frac{f(a) - 1}{a - 1} \right] (\varphi_a, \varphi)$$

and the result follows as allowing $a \rightarrow 1$. $\square$

Our main results are the following Theorem and Corollary which result from the application of Lemma 2.1 to the Brown-Ravenhall operator $\mathbf{b}$.

Theorem 2.2. *Let $Z < Z_c$ and let $\mathbf{b}$ be defined by (1.11) in $L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ with form domain $\mathcal{D}(\beta) = L^2(\mathbb{R}^3; \sqrt{1 + p^2} d\mathbf{p}) \otimes \mathbb{C}^2$. If λ is an eigenvalue of $\mathbf{b}$ with eigenvector φ , then*

$$\begin{aligned} \lambda \|\varphi\|^2 &= \int_{\mathbb{R}^3} \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)} |\varphi(\mathbf{p})|^2 d\mathbf{p} \\ &\quad - \frac{\gamma}{2} \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^1(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left[\frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} + \frac{1}{\mathbf{e}(p')} - \frac{\mathbf{e}(0)}{\mathbf{e}(p')^2} \right] d\mathbf{p}' d\mathbf{p} \\ &\quad + \frac{\gamma}{2} \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left[\frac{1}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} + \frac{1}{\mathbf{e}(p')} + \frac{\mathbf{e}(0)}{\mathbf{e}(p')^2} \right] d\mathbf{p}' d\mathbf{p}, \end{aligned} \quad (2.2)$$

where

$$\mathbf{K}^1(\mathbf{p}', \mathbf{p}) = \frac{[\mathbf{e}(p') + \mathbf{e}(0)][\mathbf{e}(p) + \mathbf{e}(0)] \mathbf{1}_2}{\mathbf{n}(p') |\mathbf{p} - \mathbf{p}'|^2 \mathbf{n}(p)}, \quad (2.3)$$

$$\mathbf{K}^2(\mathbf{p}', \mathbf{p}) = \frac{c^2(\mathbf{p}' \cdot \boldsymbol{\sigma})(\mathbf{p} \cdot \boldsymbol{\sigma})}{\mathbf{n}(p') |\mathbf{p} - \mathbf{p}'|^2 \mathbf{n}(p)}. \quad (2.4)$$

Corollary 2.3. *Let $Z < Z_c$ and $\mathbf{b}\varphi = \lambda\varphi$. Then*

$$\begin{aligned} &\left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} d\mathbf{p} \\ &\quad = \gamma \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right] d\mathbf{p}' d\mathbf{p} \\ &\quad \quad - \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} d\mathbf{p}. \end{aligned} \quad (2.5)$$

Before proving Theorem 2.2 and Corollary 2.3, we need the following lemma. We shall hereafter in this section write $\mathbf{b}_m$, $\mathbf{e}_m$, $\mathbf{K}_m$, to indicate the dependence on m .

Lemma 2.4. *For any $m \in \mathbb{R}_+$, $\mathbf{b}_m$ and $\mathbf{b}_0$ have the same domain and $\mathbf{b}_m - \mathbf{b}_0$ can be extended to a bounded operator on $L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$.*

Proof. The result is essentially proved in [8, Theorem 1] but we give a short direct proof for completeness. The following estimates are readily verified (cf [8]):

$$0 \leq \mathbf{e}_m(p) - \mathbf{e}_0(p) \leq mc^2,$$

$$\begin{aligned} |\mathbf{K}_m(\mathbf{p}', \mathbf{p}) - \mathbf{K}_0(\mathbf{p}', \mathbf{p})| &\leq \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \left\{ \frac{m}{2\mathbf{e}_m(p)} + \frac{m}{2\mathbf{e}_m(p')} + \frac{m^2}{4\mathbf{e}_m(p)\mathbf{e}_m(p')} \right\} \\ &\leq \frac{k(m)}{|\mathbf{p} - \mathbf{p}'|^2} \left\{ \frac{1}{\mathbf{e}_m(p)} + \frac{1}{\mathbf{e}_m(p')} \right\}, \end{aligned}$$

where $k(m)$ is a constant depending on m . Hence, for $\varphi \in \mathcal{D}(\beta_m) = \mathcal{D}(\beta_0)$,

$$\begin{aligned} |\beta_m[\varphi] - \beta_0[\varphi]| &\leq mc^2 \|\varphi\|^2 \\ &\quad + \gamma k(m) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \left\{ \frac{1}{\mathbf{e}_m(p)} + \frac{1}{\mathbf{e}_m(p')} \right\} |\varphi(\mathbf{p})| |\varphi(\mathbf{p}')| d\mathbf{p}' d\mathbf{p}. \end{aligned}$$

By the Cauchy-Schwarz inequality, the last integral is no greater than

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \left\{ \frac{1}{\mathbf{e}_m(p)} + \frac{1}{\mathbf{e}_m(p')} \right\} \frac{h(\mathbf{p}')}{h(\mathbf{p})} d\mathbf{p}' d\mathbf{p},$$

where we choose $h(\mathbf{p}) = |\mathbf{p}|^{-\alpha}$, $1 < \alpha < 2$. Then (see [6, p124])

$$\begin{aligned} \int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{h(\mathbf{p}')}{h(\mathbf{p})\mathbf{e}_m(p)} d\mathbf{p}' &= \frac{|\mathbf{p}|^\alpha}{\mathbf{e}_m(p)} \int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{1}{|\mathbf{p}'|^\alpha} d\mathbf{p}' \\ &\leq \frac{|\mathbf{p}|^{\alpha-1}}{c} \int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{1}{|\mathbf{p}'|^\alpha} d\mathbf{p}' = O(1) \quad \text{if } 1 < \alpha < 3, \end{aligned}$$

$$\int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{h(\mathbf{p}')}{h(\mathbf{p})\mathbf{e}_m(p')} d\mathbf{p}' \leq \frac{|\mathbf{p}|^\alpha}{c} \int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{1}{|\mathbf{p}'|^{\alpha+1}} d\mathbf{p}' = O(1) \quad \text{if } 0 < \alpha < 2.$$

Hence,

$$|\beta_m[\varphi] - \beta_0[\varphi]| \leq k(m) \|\varphi\|^2$$

and $(\beta_m - \beta_0)[\cdot, \cdot]$ can be extended to a bounded sesquilinear form on $\left\{ L^2(\mathbb{R}^3) \otimes \mathbb{C}^2 \right\} \times \left\{ L^2(\mathbb{R}^3) \otimes \mathbb{C}^2 \right\}$. It follows that $\mathbf{b}_m$ and $\mathbf{b}_0$ have the same domain and this in turn implies the rest of the lemma. $\square$

Proof of Theorem 2.2. We apply Lemma 2.1 to $T = \mathbf{b}_m$, $f(a) = a$ and $U(a)$ defined by $U(a)\varphi(\mathbf{p}) = a^{-3/2}\varphi(\mathbf{p}/a) =: \varphi_a(\mathbf{p})$. Then $U(a) \rightarrow \mathbf{I}$ strongly in $L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ and

$$T_a := aU(a)\mathbf{b}_mU(a)^{-1} = \mathbf{b}_{ma}.$$

By Lemma 2.4, an eigenvector φ of $\mathbf{b}_m$, lies in $\mathcal{D}(\mathbf{b}_{ma})$ for any a . Thus, it remains to evaluate the limit on the left-hand side of (2.1). We have

$$\begin{aligned} \left(\varphi_a, \left[\frac{\mathbf{b}_{ma} - \mathbf{b}_m}{a-1} \right] \varphi \right) &= \int_{\mathbb{R}^3} \left[\frac{\mathbf{e}_{ma}(\mathbf{p}) - \mathbf{e}_m(\mathbf{p})}{a-1} \right] |\varphi(\mathbf{p})|^2 d\mathbf{p} \\ &\quad - \gamma \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi_a(\mathbf{p}')^* \left[\frac{\mathbf{K}_{ma}(\mathbf{p}', \mathbf{p}) - \mathbf{K}_m(\mathbf{p}', \mathbf{p})}{a-1} \right] \varphi(\mathbf{p}) d\mathbf{p}' d\mathbf{p} =: I_1 - \gamma I_2, \end{aligned} \quad (2.6)$$

and (2.2) will follow if we can justify taking the limit as $a \rightarrow 1$ under the integral signs on the right-hand side of (2.6).

In I_1 the integrand is majorised by $|\varphi(\mathbf{p})|^2$ and so the dominated convergence theorem applies. We write I_2 as

$$\begin{aligned} &\iint_{\mathbb{R}^3 \times \mathbb{R}^3} [\varphi_a(\mathbf{p}') - \varphi(\mathbf{p}')]^* \left[\frac{\mathbf{K}_{ma}(\mathbf{p}', \mathbf{p}) - \mathbf{K}_m(\mathbf{p}', \mathbf{p})}{a-1} \right] \varphi(\mathbf{p}) d\mathbf{p}' d\mathbf{p} \\ &+ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi(\mathbf{p}')^* \left[\frac{\mathbf{K}_{ma}(\mathbf{p}', \mathbf{p}) - \mathbf{K}_m(\mathbf{p}', \mathbf{p})}{a-1} \right] \varphi(\mathbf{p}) d\mathbf{p}' d\mathbf{p} =: I_3 + I_4, \end{aligned}$$

and use the readily verified estimate

$$\left| \frac{\mathbf{K}_{ma}(\mathbf{p}', \mathbf{p}) - \mathbf{K}_m(\mathbf{p}', \mathbf{p})}{a-1} \right| \leq k_a / |\mathbf{p} - \mathbf{p}'|^2,$$

where $k_a \rightarrow 1$ as $a \rightarrow 1$. Thus I_3 bounded by

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |\varphi_a(\mathbf{p}') - \varphi(\mathbf{p}')| |\varphi(\mathbf{p})| \frac{d\mathbf{p}' d\mathbf{p}}{|\mathbf{p} - \mathbf{p}'|^2}$$

and, on using the Parseval identity and $\mathcal{F}(\frac{1}{|\cdot|}) = \sqrt{\frac{2}{\pi}} \frac{1}{|\cdot|^2}$ (with $\mathcal{F}$ now the standard Fourier transform with $\hbar = 1$), this is equal to

$$2\pi^2 \int_{\mathbb{R}^3} u(\mathbf{x}) v(\mathbf{x}) \frac{d\mathbf{x}}{|\mathbf{x}|},$$

where $\hat{u}(\mathbf{p}) = |\varphi_a(\mathbf{p}) - \varphi(\mathbf{p})|$ and $\hat{v}(\mathbf{p}) = |\varphi(\mathbf{p})|$,

$$\leq \pi^3 \left(\int_{\mathbb{R}^3} |\varphi_a(\mathbf{p}) - \varphi(\mathbf{p})|^2 |\mathbf{p}| d\mathbf{p} \right)^{1/2} \left(\int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 |\mathbf{p}| d\mathbf{p} \right)^{1/2}$$

by the Cauchy-Schwarz inequality and Kato's inequality [5, p. 307]. Thus $I_3 \rightarrow 0$ as $a \rightarrow 1$ since $\varphi \in L^2(\mathbb{R}^3; \sqrt{1+p^2} d\mathbf{p}) \otimes \mathbb{C}^2$. In I_4 the integrand is majorised by the function $\frac{1}{|\mathbf{p}-\mathbf{p}'|^2} |\varphi(\mathbf{p}')| |\varphi(\mathbf{p})|$ which is integrable by Kato's inequality and hence the dominated convergence theorem applies. $\square$

Proof of Corollary 2.3. From (2.2) we have

$$\begin{aligned} \lambda \|\varphi\|^2 &= \mathbf{e}(0)^2 \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{d\mathbf{p}}{\mathbf{e}(p)} \\ &\quad - \gamma \mathbf{e}(0) \operatorname{Re} \left[\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left\{ \frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} \right\} d\mathbf{p}' d\mathbf{p} \right] \\ &\quad + \gamma \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left\{ \frac{1}{\mathbf{e}(p)} + \frac{1}{\mathbf{e}(p')} \right\} d\mathbf{p}' d\mathbf{p}. \end{aligned} \quad (2.7)$$

Also, for all $\psi \in \mathcal{D}(\beta)$,

$$\begin{aligned} \int_{\mathbb{R}^3} \mathbf{e}(p) \psi^*(\mathbf{p}) \varphi(\mathbf{p}) d\mathbf{p} - \gamma \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \psi^*(\mathbf{p}') \mathbf{K}(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) d\mathbf{p}' d\mathbf{p} \\ = \lambda \int_{\mathbb{R}^3} \psi^*(\mathbf{p}) \varphi(\mathbf{p}) d\mathbf{p}. \end{aligned} \quad (2.8)$$

We choose $\psi(\mathbf{p}) = \left[\frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} \right] \varphi(\mathbf{p})$ in (2.8): clearly $\psi \in \mathcal{D}(\beta)$. Then

$$\begin{aligned} \gamma \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left\{ \frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} \right\} d\mathbf{p}' d\mathbf{p} \\ = \int_{\mathbb{R}^3} [\mathbf{e}(p) - \lambda] \left\{ \frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} \right\} |\varphi(\mathbf{p})|^2 d\mathbf{p}. \end{aligned} \quad (2.9)$$

On substituting (2.9) in (2.7),

$$\begin{aligned} \lambda \|\varphi\|^2 &= \mathbf{e}(0)^2 \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{d\mathbf{p}}{\mathbf{e}(p)} - \mathbf{e}(0) \int_{\mathbb{R}^3} [\mathbf{e}(p) - \lambda] \left\{ \frac{1}{\mathbf{e}(p)} - \frac{\mathbf{e}(0)}{\mathbf{e}(p)^2} \right\} |\varphi(\mathbf{p})|^2 d\mathbf{p} \\ &\quad + \gamma \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left\{ \frac{1}{\mathbf{e}(p)} + \frac{1}{\mathbf{e}(p')} \right\} d\mathbf{p}' d\mathbf{p}, \end{aligned}$$

whence

$$\begin{aligned} \lambda \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} d\mathbf{p} &= \mathbf{e}(0) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ \frac{\mathbf{e}(0)}{\mathbf{e}(p)} - 1 + \frac{\mathbf{e}(0)}{\mathbf{e}(p)} \right\} d\mathbf{p} \\ &\quad + \gamma \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right] d\mathbf{p}' d\mathbf{p}, \end{aligned}$$

and

$$\begin{aligned}
& (\lambda - \mathbf{e}(0)) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} d\mathbf{p} \\
&= \mathbf{e}(0) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][\mathbf{e}(0) - 2\mathbf{e}(p)]}{\mathbf{e}(p)^2} d\mathbf{p} \\
&+ \gamma \mathbf{e}(0) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{K}^2(\mathbf{p}', \mathbf{p}) \varphi(\mathbf{p}) \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right] d\mathbf{p}' d\mathbf{p}. \quad (2.10)
\end{aligned}$$

The corollary is therefore proved. $\square$

3. THE ABSENCE OF EMBEDDED EIGENVALUES

Theorem 3.1. *If $Z < Z_c$, the operator $\mathbf{b}$ has no eigenvalues in $[\max\{1, 2\alpha Z - 1/2\}mc^2, \infty)$. In particular, if $Z \leq Z'_c = 3/(4\alpha)$, there are no eigenvalues in $[mc^2, \infty)$ and the spectrum of $\mathbf{b}$ is absolutely continuous.*

Proof. From (2.5)

$$\begin{aligned}
& \left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} d\mathbf{p} \\
& \leq \gamma \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \varphi^*(\mathbf{p}') \mathbf{H}(\mathbf{p}', \mathbf{p}) |\varphi(\mathbf{p})| |\varphi(\mathbf{p}')| d\mathbf{p}' d\mathbf{p} \\
& \quad - \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} d\mathbf{p}, \quad (3.1)
\end{aligned}$$

where

$$\mathbf{H}(\mathbf{p}', \mathbf{p}) = \frac{c^2 p p'}{\mathbf{n}(p') \mathbf{n}(p) |\mathbf{p} - \mathbf{p}'|^2} \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right].$$

By the Cauchy-Schwarz inequality

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \mathbf{H}(\mathbf{p}', \mathbf{p}) |\varphi(\mathbf{p}')| |\varphi(\mathbf{p})| d\mathbf{p}' d\mathbf{p} \leq c^2 \int_{\mathbb{R}^3} \frac{p^2}{\mathbf{n}^2(p)} |\varphi(\mathbf{p})|^2 d\mathbf{p} \int_{\mathbb{R}^3} \frac{h(\mathbf{p}')}{h(\mathbf{p})} \mathbf{J}(\mathbf{p}', \mathbf{p}) d\mathbf{p}', \quad (3.2)$$

where

$$\mathbf{J}(\mathbf{p}', \mathbf{p}) = \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right]$$

and $h(\cdot)$ is an arbitrary positive function. We make the choice $h(\mathbf{p}) = p^{-3/2}$ and use the following (see [6, p 124]):

$$\int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{1}{|\mathbf{p}'|^{3/2}} d\mathbf{p}' = 4\pi^2 p^{-1/2},$$

$$\int_{\mathbb{R}^3} \frac{1}{|\mathbf{p} - \mathbf{p}'|^2} \frac{1}{|\mathbf{p}'|^{5/2}} d\mathbf{p}' = 4\pi^2 p^{-3/2}.$$

On substituting in (3.2), we have

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \mathbf{H}(\mathbf{p}', \mathbf{p}) |\varphi(\mathbf{p}')| |\varphi(\mathbf{p})| d\mathbf{p}' d\mathbf{p} < 4\pi^2 c^2 \int_{\mathbb{R}^3} \frac{p^2}{\mathbf{n}^2(p)} \left[\frac{p}{\mathbf{e}(p)} + \frac{1}{c} \right] |\varphi(\mathbf{p})|^2 d\mathbf{p}$$

and hence (3.1) yields

$$\begin{aligned} & \left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} d\mathbf{p} \\ & < \int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 \left\{ 4\pi^2 c^2 \gamma \frac{p^2}{\mathbf{n}^2(p)} \left[\frac{p}{\mathbf{e}(p)} + \frac{1}{c} \right] - \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} \right\} d\mathbf{p}. \end{aligned}$$

On replacing $\mathbf{p}$ by $m c \mathbf{p}$ and simplifying, we derive

$$0 < \int_{\mathbb{R}^3} |\varphi(m c \mathbf{p})|^2 \left(1 - \frac{1}{\sqrt{p^2 + 1}} + \frac{1}{p^2 + 1} \right) \left(1 - \frac{\lambda}{m c^2} + \Phi(p) \right) d\mathbf{p}, \quad (3.3)$$

where

$$\begin{aligned} \Phi(p) &= \frac{p^2(p + \sqrt{p^2 + 1})}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ 2\pi^2 \left(\frac{\gamma}{c} \right) - \frac{2\sqrt{p^2 + 1} - 1}{p + \sqrt{p^2 + 1}} \right\} \\ &\leq \frac{p^2(p + \sqrt{p^2 + 1})}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ 2\pi^2 \left(\frac{\gamma}{c} \right) - \frac{3}{4} \right\} \end{aligned}$$

since $\min_{\mathbb{R}_+} \left\{ \frac{2\sqrt{p^2+1}-1}{p+\sqrt{p^2+1}} \right\} = \frac{3}{4}$. Thus, if $2\pi^2(\frac{\gamma}{c}) \equiv \alpha Z \leq \frac{3}{4}$, (3.3) implies that $\lambda < m c^2$. Since $\sup_{\mathbb{R}_+} \left\{ \frac{p^2(p+\sqrt{p^2+1})}{(\sqrt{p^2+1}+1)(p^2+2-\sqrt{p^2+1})} \right\} = 2$, it follows from (3.3) that, if $2\pi^2(\frac{\gamma}{c}) \equiv \alpha Z > \frac{3}{4}$, $\lambda < m c^2 [1 + 2\alpha Z - \frac{3}{2}]$. The theorem is therefore proved $\square$

Finally, we analyse the implications of the virial theorem for the operators $b_{l,s}$.

In the partial wave decomposition of $\mathbf{B}$ and $\mathbf{b}$ (see [3]), spinors $\varphi \in L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ are expanded in terms of spherical spinors $\Omega_{l,n,s}$

$$\varphi(\mathbf{p}) = \sum_{(l,n,s) \in I} p^{-1} \varphi_{l,n,s}(p) \Omega_{l,n,s}(\omega) \quad (3.4)$$

say, where $I = \{(l, n, s) : l \in \mathcal{N}_0, n = -l - \frac{1}{2}, \dots, l + \frac{1}{2}, s = \pm \frac{1}{2}, \Omega_{l,n,s} \neq 0\}$ and

$$\int_{\mathbb{R}^3} |\varphi(\mathbf{p})|^2 d\mathbf{p} = \sum_{(l,n,s) \in I} \int_0^\infty |\varphi_{l,n,s}(p)|^2 dp. \quad (3.5)$$

On substituting in (1.5), we get

$$\beta[\varphi, \psi] = \sum_{(l,n,s) \in I} \beta_{l,s}[\varphi_{l,n,s}, \psi_{l,n,s}], \quad (3.6)$$

where

$$\begin{aligned} \beta_{l,s}[\varphi_{l,n,s}, \psi_{l,n,s}] &= \int_0^\infty \mathbf{e}(p) \overline{\varphi}_{l,n,s}(p) \psi_{l,n,s}(p) \, dp \\ &\quad - \frac{\alpha c Z}{\pi} \int_0^\infty \int_0^\infty \overline{\varphi}_{l,n,s}(p') k_{l,s}(p', p) \psi_{l,n,s}(p) \, dp' \, dp \quad (3.7) \end{aligned}$$

and, with Q_l denoting Legendre functions of the second kind,

$$\begin{aligned} k_{l,s}(p', p) &= \frac{[\mathbf{e}(p') + \mathbf{e}(0)] Q_l\left(\frac{1}{2}\left[\frac{p'}{p} + \frac{p}{p'}\right]\right) [\mathbf{e}(p) + \mathbf{e}(0)]}{\mathbf{n}(p') \mathbf{n}(p)} + \frac{c^2 p' Q_{l+2s}\left(\frac{1}{2}\left[\frac{p'}{p} + \frac{p}{p'}\right]\right) p}{\mathbf{n}(p') \mathbf{n}(p)} \\ &=: k_{l,m}^1(p', p) + k_{l,m}^2(p', p) \quad (3.8) \end{aligned}$$

say. For $Z < Z_c$, the forms with domain $L^2(0, \infty; \sqrt{1+p^2} \, dp)$ are closed and positive in $L^2(0, \infty)$, and we shall denote the associated self-adjoint operators by $b_{l,s}$. These operators $b_{l,s}$ coincide with the operators $b_{l,s}$ of Tix in [8] when the later are self-adjoint, but are otherwise their Friedrichs extensions. Note that it follows from [3, Theorem2] that for all values of l, s , $\sigma_{\text{ess}}(b_{l,s}) = [mc^2, \infty)$.

Theorem 3.2. *For $(l, s) \neq (1, -\frac{1}{2})$, the operators $b_{l,s}$ have no eigenvalues in $[mc^2, \infty)$ if $Z < Z_c$. If $(l, s) = (1, -\frac{1}{2})$, $b_{l,s}$ has no eigenvalues in $[\max\{1, 2\alpha Z - 1/2\} mc^2, \infty)$: in particular $b_{1,-1/2}$ has no eigenvalues in $[mc^2, \infty)$ if $Z \leq Z'_c = \frac{3}{4\alpha}$.*

Proof. The analogue of (2.5) is

$$\begin{aligned} &\left(\frac{\lambda}{\mathbf{e}(0)} - 1\right) \int_0^\infty |\varphi_{l,n,s}(p)|^2 \left\{1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2}\right\} \, dp \\ &= 2\pi\gamma \int_0^\infty \int_0^\infty \overline{\varphi}_{l,n,s}(p') \mathbf{k}_{l,s}^2(p', p) \varphi(p) \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)}\right] \, dp' \, dp \\ &\quad - \int_0^\infty |\varphi_{l,n,s}(p)|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} \, dp. \quad (3.9) \end{aligned}$$

Since $Q_0(t) \geq \dots \geq Q_l(t) \geq 0$ for all $t > 1$, it follows that for all $(l, s) \neq (1, -1/2)$

$$k_{l,s}^2(p', p) \leq \frac{c^2 p' p}{\mathbf{n}(p') \mathbf{n}(p)} Q_1\left(\frac{1}{2}\left[\frac{p'}{p} + \frac{p}{p'}\right]\right)$$

and hence, on writing φ for $\varphi_{l,n,s}$ in (3.9), we get

$$\begin{aligned} & \left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_0^\infty |\varphi(p)|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} \mathrm{d}p \\ & \leq 2\pi\gamma c^2 \int_0^\infty |\varphi(p)|^2 \frac{p^2}{\mathbf{n}^2(p)h(p)} \mathrm{d}p \int_0^\infty h(p') Q_1 \left(\frac{1}{2} \left[\frac{p'}{p} + \frac{p}{p'} \right] \right) \left(\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right) \mathrm{d}p' \\ & \quad - \int_0^\infty |\varphi(p)|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} \mathrm{d}p, \quad (3.10) \end{aligned}$$

for any positive function $h(\cdot)$, on applying the Cauchy-Schwarz inequality. We choose $h(t) = 1/t$, so that in the first term on the right-hand side of (3.10) we have

$$\begin{aligned} & 2\pi\gamma c^2 \int_0^\infty |\varphi(p)|^2 \frac{p^2}{\mathbf{n}^2(p)} \left\{ \frac{p}{\mathbf{e}(p)} \int_0^\infty \frac{1}{p'} Q_1 \left(\frac{1}{2} \left[\frac{p'}{p} + \frac{p}{p'} \right] \right) \mathrm{d}p' \right. \\ & \quad \left. + p \int_0^\infty \frac{1}{p' \mathbf{e}(p')} Q_1 \left(\frac{1}{2} \left[\frac{p'}{p} + \frac{p}{p'} \right] \right) \mathrm{d}p' \right\} \mathrm{d}p \\ & = 2\pi\gamma c^2 \int_0^\infty |\varphi(p)|^2 \frac{p^2}{\mathbf{n}^2(p)} \left\{ I_1 + I_2 \right\} \mathrm{d}p \end{aligned}$$

say. Let $g_1(u) = Q_1 \left(\frac{1}{2} \left[u + \frac{1}{u} \right] \right)$. Then

$$cI_1 \leq \int_0^\infty g_1(u) \frac{\mathrm{d}u}{u} = 2,$$

(see [2, (3.8)] or [3, § 2.3]), and

$$cI_2 \leq \int_0^\infty g_1(u) \frac{\mathrm{d}u}{u^2} = \int_0^\infty g_1(u) \mathrm{d}u.$$

We have (see [2, (3.7)])

$$\begin{aligned} \int_0^1 g_1(u) \mathrm{d}u &= \frac{1}{2} \int_0^1 u \ln \left(\frac{u+1}{1-u} \right) \mathrm{d}u + \frac{1}{2} \int_0^1 \frac{1}{u} \ln \left(\frac{u+1}{1-u} \right) \mathrm{d}u - \int_0^1 \mathrm{d}u \\ &= \frac{1}{2} + \frac{1}{2} \frac{\pi^2}{4} - 1 = \frac{\pi^2}{8} - \frac{1}{2} \end{aligned}$$

and

$$\begin{aligned} \int_1^\infty g_1(u) \, du &= \frac{1}{2} \int_1^\infty \frac{1}{u} \ln\left(\frac{u+1}{u-1}\right) \, du + \int_1^\infty \left\{ \frac{u}{2} \ln\left(\frac{u+1}{u-1}\right) - 1 \right\} \, du \\ &= \frac{\pi^2}{8} + \left[\frac{1}{4}(u^2 - 1) \ln\left(\frac{u+1}{u-1}\right) - \frac{u}{2} \right]_1^\infty = \frac{\pi^2}{8} + \frac{1}{2}. \end{aligned}$$

Hence $cI_2 \leq \frac{\pi^2}{4}$, and from (3.10)

$$\begin{aligned} &\left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_0^\infty |\varphi(p)|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} \, dp \\ &\leq 2\pi\gamma c \int_0^\infty |\varphi(p)|^2 \frac{p^2}{\mathbf{n}^2(p)} \left(\frac{\pi^2}{4} + 2 \right) \, dp - \int_0^\infty |\varphi(p)|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} \, dp. \end{aligned}$$

On replacing p by mcp and simplifying, we obtain

$$0 < \int_0^\infty |\varphi(mcp)|^2 \left(1 - \frac{1}{\sqrt{p^2 + 1}} + \frac{1}{p^2 + 1} \right) \left(1 - \frac{\lambda}{mc^2} + \Psi(p) \right) \, dp,$$

where

$$\begin{aligned} \Psi(p) &= \frac{p^2(\sqrt{p^2 + 1})}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ \pi \left(\frac{\gamma}{c} \right) \left(\frac{\pi^2}{4} + 2 \right) - \frac{2\sqrt{p^2 + 1} - 1}{\sqrt{p^2 + 1}} \right\} \\ &\leq \frac{p^2\sqrt{p^2 + 1}}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ \frac{2}{\pi^2 + 4} \left(\frac{\pi^2}{4} + 2 \right) - 1 \right\} < 0 \end{aligned}$$

for $2\pi^2 \frac{\gamma}{c} \equiv \alpha Z \leq \alpha Z_c$. Hence $\lambda < mc^2$.

For the case $(l, s) = (1, -1/2)$, the Legendre function Q_1 in (3.10) has to be replaced by Q_0 , and we have to consider

$$\frac{1}{h(p)} \int_0^\infty h(p') Q_0 \left(\frac{1}{2} \left[\frac{p'}{p} + \frac{p}{p'} \right] \right) \left[\frac{1}{\mathbf{e}(p')} + \frac{1}{\mathbf{e}(p)} \right] \, dp'. \quad (3.11)$$

We now make the choice $h(u) = 1/\sqrt{u}$, and, with $g_0(u) = Q_0 \left(\frac{1}{2} \left[u + \frac{1}{u} \right] \right) \equiv \ln \left| \frac{u+1}{u-1} \right|$, (3.11) becomes $J_1 + J_2$ say, where

$$J_1 = p \int_0^\infty \frac{1}{\sqrt{u}\mathbf{e}(pu)} g_0(u) \, du < \frac{1}{c} \int_0^\infty \frac{g_0(u)}{u^{3/2}} \, du$$

and

$$J_2 = \frac{p}{\mathbf{e}(p)} \int_0^\infty \frac{1}{\sqrt{u}} g_0(u) \, du.$$

Since

$$\begin{aligned}\int_0^1 \frac{g_0(u)}{u^{1/2}} du &= \int_1^\infty \frac{g_0(u)}{u^{3/2}} du, \\ \int_1^\infty \frac{g_0(u)}{u^{1/2}} du &= \int_0^1 \frac{g_0(u)}{u^{3/2}} du\end{aligned}$$

we have

$$\begin{aligned}\int_0^\infty \frac{g_0(u)}{u^{1/2}} du &= \int_0^\infty \frac{g_0(u)}{u^{3/2}} du = \int_1^\infty g_0(u) \left(1 + \frac{1}{u}\right) \frac{du}{\sqrt{u}} \\ &= 4 \int_1^\infty (\ln y) \frac{y}{(y^2 - 1)^{3/2}} dy\end{aligned}$$

on setting $\frac{u+1}{u-1} = y$,

$$= \int_1^\infty (\ln z) \frac{dz}{(z-1)^{3/2}} = 2 \int_1^\infty \frac{1}{z(z-1)^{1/2}} dz = 2\pi.$$

Hence, on substituting in (3.10) (with Q_1 replaced by Q_0) we infer that

$$\begin{aligned}& \left(\frac{\lambda}{\mathbf{e}(0)} - 1 \right) \int_0^\infty |\varphi(p)|^2 \left\{ 1 - \frac{\mathbf{e}(0)}{\mathbf{e}(p)} + \frac{\mathbf{e}(0)^2}{\mathbf{e}(p)^2} \right\} dp \\ & < 2\pi\gamma c \int_0^\infty |\varphi(p)|^2 \frac{p^2}{\mathbf{n}^2(p)} \left[2\pi \left(\frac{cp}{\mathbf{e}(p)} + 1 \right) \right] dp - \int_0^\infty |\varphi(p)|^2 \frac{[\mathbf{e}(p) - \mathbf{e}(0)][2\mathbf{e}(p) - \mathbf{e}(0)]}{\mathbf{e}(p)^2} dp.\end{aligned}$$

On replacing p by mcp and simplifying, this yields

$$0 < \int_0^\infty |\varphi(mcp)|^2 \left(1 - \frac{1}{\sqrt{p^2 + 1}} + \frac{1}{p^2 + 1} \right) \left(1 - \frac{\lambda}{mc^2} + \Theta(p) \right) dp,$$

where

$$\begin{aligned}\Theta(p) &= \frac{p^2(p + \sqrt{p^2 + 1})}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ 2\pi^2 \left(\frac{\gamma}{c} \right) - \frac{2\sqrt{p^2 + 1} - 1}{p + \sqrt{p^2 + 1}} \right\} \\ &\leq \frac{p^2(\sqrt{p^2 + 1} + p)}{(\sqrt{p^2 + 1} + 1)(p^2 + 2 - \sqrt{p^2 + 1})} \left\{ 2\pi^2 \left(\frac{\gamma}{c} \right) - \frac{3}{4} \right\} \leq 0\end{aligned}$$

if $2\pi^2\left(\frac{\gamma}{c}\right) = \alpha Z \leq \alpha Z'_c = 3/4$. Hence in this case $\lambda < mc^2$. If $\alpha Z > 3/4$, we have

$$\Theta(p) \leq 2\left(\alpha Z - \frac{3}{4}\right)$$

and $\lambda < mc^2\left(\alpha Z - \frac{1}{2}\right)$. The theorem is therefore proved. $\square$

ACKNOWLEDGEMENTS

The authors are grateful to the European Union for support under the TMR grant FMRX-CT 96-0001.

REFERENCES

- [1] G.E. Brown and D.G. Ravenhall, *On the interaction of two electrons*, Proc. Roy. Soc. London A **208** (1952), pp. 552-559.
- [2] V.I. Burenkov and W.D. Evans, *On the evaluation of the norm of an integral operator associated with the stability of one-electron atoms*, to appear in Proc. Roy. Soc. Edinburgh.
- [3] William Desmond Evans, Peter Perry and Heinz Siedentop, *The spectrum of relativistic one-electron atoms according to Bethe and Salpeter*, Commun. Math. Phys. **178** (1996), pp. 733-746.
- [4] G. Hardekopf and J. Sucher, *Critical coupling constants for relativistic wave equations and vacuum breakdown in quantum electrodynamics*, Phys. Rev. A, **31**(4) (April 1985), pp. 2020-2029.
- [5] T. Kato, *Perturbation Theory for Linear Operators*, Grundlehren der mathematischen Wissenschaften, vol. 132, Springer-Verlag, Berlin, Heidelberg, New York, Second Edition, 1976.
- [6] E.H. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics, vol. 14, American Mathematical Society, 1997.
- [7] C. Tix, *Strict positivity of a relativistic Hamiltonian due to Brown and Ravenhall*, Preprint mp-arc/96-660.
- [8] C. Tix, *Self-adjointness and spectral properties of a pseudo-relativistic Hamiltonian due to Brown and Ravenhall*, Preprint mp-arc/97-441.

SCHOOL OF MATHEMATICS, UNIVERSITY OF WALES, CARDIFF, 23 SENGHENNYDD ROAD, P O BOX 926, CARDIFF CF2 4YH, GREAT BRITAIN

E-mail address, A. A. Balinsky: BalinskyA@cardiff.ac.uk

E-mail address, W. D. Evans: EvansWD@cardiff.ac.uk